

## The immunosuppressive impact of PRRS virus on the immune response following anti - erysipelas vaccination in swine from various farms

### Influența imunosupresoare a virusului PRRS asupra răspunsului imun în urma vaccinării antirujetice la porcine din diferite sisteme de exploatație

Viorica Chiurciu, C Costin Stoica, Maria Mioara Răduță, Cătălin Tudoran,  
Lucia Diaconu, Ion Iacob  
SC Romvac Company SA, Voluntari, Romania,

Correspondence: [chiurciu@romvac.ro](mailto:chiurciu@romvac.ro)

**Key words:** PRRS, rujet, vaccination, immunosuppression

**Cuvinte cheie:** PRRS, rujet, vaccinare, imunosupresie

#### Abstract

PRRS virus, the etiologic agent of Porcine Reproductive and Respiratory Syndrome by immunosuppressive action can significantly affect the immune response after vaccination. It was intended the following of the immunological reaction induced by the *Erysipelothrix rhusopathiae* vaccination from pigs from intensive system and from households. The biological material studied was provided from clinically healthy pigs of different ages. The animals were from four different locations. Serological examinations were performed by blood sampling [gathered from the confluence of jugular vein] before and after the vaccination. The investigations were performed by ELISA method. In the industrial breeding system, seroprevalence of anti PRRS presented high levels, in contrast to the low level of postvaccinal *E. rhusopathiae* antibodies. In households the incidence of PRRS virus was low and the seroconversion after the vaccination was raised. The morphopathological and bacteriological examinations performed from the lesions in various organs [lungs, lymph nodes, liver, spleen and intestine] has revealed the presence of germ association, pathogenic or potentially pathogenic. The results point the link between the existence of PRRS virus in the swine populations and post-vaccinal response, its presence interfering significantly with the vaccination protocols efficacy.

#### Rezumat

Virusul PRRS, agentul etiologic al Sindromului Respirator și Reproductiv Porcin, prin acțiunea sa imunosupresoare poate afecta semnificativ răspunsul imun postvaccinal. S-a urmărit reacția imunologică indusă de vaccinarea antirujetică la porci crescuți atât în sistem intensiv cât și în gospodăriile populației. Materialul biologic luat în studiu a fost reprezentat de suine clinic sănătoase, de diferite vârste. Animalele au provenit din 4 locații diferite. Au fost efectuate examene serologice prin prelevare de sange [din confluența jugulară] înainte și după vaccinarea antirujetică. Investigațiile au fost realizate prin metoda ELISA. În sistemul de creștere industrial, seroprevalența anticorpilor anti PRRS a prezentat niveluri ridicate, în contrast cu nivelul scăzut al anticorpilor antirujetici postvaccinali. În gospodăriile populației incidența virusului PRRS a fost scăzută iar seroconversia în urma vaccinării antirujetice a fost ridicată. În urma examenului anatomopatologic și a examenelor bacteriologice efectuate din leziuni de la nivelul diferitelor organe [pulmoni, limfonoduli, ficat, splină, intestin] s-a evidențiat prezența unor germeni de asociație, patogeni sau potențial patogeni. Rezultatele indică o legătură între existența virusului PRRS în populațiile de porcine și răspunsul imun postvaccinal, prezența acestuia putând interfeera semnificativ cu eficiența protoalelor de vaccinare.

#### Introduction

Porcine Respiratory and Reproductive Syndrome (PRRS) is a disease which occurs

both in domestic and wild swine [6], covering all categories of age, mostly sows and suckling pigs.

The disease is spread worldwide [2, 4, 8, 11].

The disease is transmitted by aerogenic, genital and transplacental routes. The spread is fast and covers all suckling pigs within 1-2 weeks only.

The etiologic agent is a 45-55 nm diameter DNA virus of family *Arteriviridae*, genus *Arterivirus*.

Virus replication takes place initially in the alveolar macrophages, the impairment of which shall enhance secondary infections with bacteria such as *Actinobacillus pleuropneumoniae*, *Haemophilus parasuis*, *Pasteurella multocida* etc.

It causes reproductive disorders, abortions or mummified fetuses. Piglets infected during pregnancy develop growth disorders, shortness of breath, anorexia, fever and horripilation.

The pathological examination may reveal interstitial pneumonia [1, 3, 5, 7].

A previous study conducted by our research team in 2 farms and households demonstrated that PRRS seroprevalence was 69.5% (313 positive samples of 450) and 0% in households [10].

Another study conducted by our research team on forest swine, using serological tests revealed no PRRS in boars [9].

This paper describes a study carried out in swine farms and households from our country, being aware of the existence of global pathogenic agents known for their immunosuppressive effect on swine.

They can weaken the defense capacity of the organism resulting in increased risk of occurrence and severe progress of diseases.

Such a pathogen which can significantly interfere with the efficacy of vaccination protocols is the PRRS virus, the etiologic agent of Porcine Respiratory and Reproductive Syndrome.

For this reason, the immunological reaction induced by anti-erysipelas vaccination has been observed both in farm and household - reared swine, taking as reference the impact of PRRS virus on the anti-erysipelas immune response since anti-erysipelas vaccination is a foreseen action carried out in the current technology applied in swine farms.

## Materials and methods

The biological material under study consisted of clinically healthy swine of various ages.

The animals originated from 4 distinct swine-rearing facilities (A, B, C, D): 3 intensive system farms (Farm A: 172 animals, Farm B: 90 animals, Farm C: 99 animals) and from households (50 animals).

Serological examinations were carried out by blood sampling (from the jugular) before and after anti-erysipelas vaccination.

Several types of live and subunitary anti-erysipelas vaccine were used from various manufacturers.

Using ELISA technique, 728 blood serum samples were analyzed from 512 swine: 390 young pigs (1.5 – 4 months old), 70 fat pigs (6 months – 1 year old) and 52 sows in waiting period. PRRS-specific Ig G antibodies and *Erysipelothrix rhusiopathiae* from blood serum were detected using ELISA kit for PRRS (HerdCheck-PRRS X3 IDEXX Laboratories, USA) and *Erysipelothrix rhusiopathiae* (*Erysipelothrix rhusiopathiae* SE / MR, Cypress Diagnostics).

For bacteriological examinations, samples of nasal discharge were taken on sterile swabs from trial animals and wounded organs were taken from dead animals (lungs, lymph nodes, liver, spleen, intestine).

The bacteriological examinations consisted of seeding on enrichment and selective media. The identification was carried out with API multitesting systems (Biomerieux).

## Results and discussions


Farm A (Picovit Farm) relies on farm (intensive) rearing of young pigs, purchasing them from abroad (Holland) and from inside the country as well.

Pigs are purchased at the age of 65-70 days and reared up to the age of 150-160 days, when they are delivered.

Anti-PRRS and erysipelas vaccinations are performed within this interval. Serological surveillance is carried out for classic swine pest.

A total of 329 samples were tested, 188 of which were PRRS-positive (57.1%),

Seroconversion after anti-erysipelas vaccination was 5.8%, i.e. 11 positive samples of 187 tested (figure 1).




**Figure 1.** Immune response after anti-erysipelas vaccination in farm A

Farm B (Periș Farm) is a closed-circuit farm (breeding, growing and fattening) and the breeding material (gilts and young boars) is from the own breeding material.

The sanitary-veterinary surveillance is carried out for pleuropneumonia, Aujeszky disease and brucellosis and the self-control for PRRS and Circovirus is carried out by serological tests and pathological examinations as well.

Anti-erysipelas vaccination is carried out in this farm in 90-120 days old piglets and in sows 3 days after parturition and 3 days before weaning.

170 samples were tested and 161 were PRRS-positive, and seroconversion after anti-erysipelas vaccination was 23.7%, i.e. 19 positive samples of 80 tested (figure 2).



**Figure 2.** Immune response after anti-erysipelas vaccination in farm B

In farm C (PigCom Satu Nou), breeding, growing and fattening of pigs are carried out in 13 shelters (pens), accordingly:


- 3 breeding pens, exclusively for artificial fertilization, the seminal material being provided by 28 terminal boars. The seminal material has been previously imported as well, probably leading to the source of PRRS, circovirus infection;
- 2 pregnancy pens intended for parturition and growing of progenies until weaning (30 - 32 days);
- 2 youth growing pens where the most representative gilt specimens were selected from the phenotypic point of view to replace the breeding sows;
- 6 pens for fattening pigs

The female breeding material consists of the in-house youth replacing the sows, exploited 2.1 times per year and terminal seminal donors acquired from SC PIG Romania, SC Dan Bred Ltd. in Argeș County and in the future from special farms in Hungary, currently negotiating the purchase terms with.


After 30-32 days in maternity, pigs become growing youth using the "all in - all out" technique according to which a small number consists of growing animals on which a certain flora is transplanted determining a relatively high percentage of 8-10% loss by mortality, a frequent phenomenon in this category of pigs.

Fat pigs are reared on ground up to the age and weight required for delivery, when their assessment is carried out at slaughter. 99 samples were tested and 41 were PRRS-positive (49.4%).

Seroconversion after anti-erysipelas vaccination was 9%, i.e. 9 positive samples of 99 tested (figure 3).



**Figure 3.** Immune response after anti-erysipelas vaccination in farm C



**Figure 4.** Immune response after anti-erysipelas vaccination in households

130 samples from households were tested obtaining 3 PRRS-positive samples (2.3%). Seroconversion after anti-erysipelas vaccination was 57.5%, i.e. 23 positive samples of 40 tested (fig. 4).


The synthesis of serological test results regarding PRRS and immune response seroprevalence after anti-erysipelas vaccination is described in table 1.

**Table 1. PRRS virus incidence in swine and postvaccinal anti-erysipelas immune response**

Locations	IgG ELISA results			
	PRRS No. of positive samples / tested samples	Seroprevalence (%)	Erysipelas No. of positive samples / tested samples	Seroconversion (%)
FARM A	188/329	57,1	11/187	5,8
FARM B	161/170	94,7	19/80	23,7
FARM C	41/99	41,4	9/99	9
Households	3/130	2,3	23/40	57,5
<b>TOTAL</b>	<b>393/728</b>	<b>54</b>	<b>62/406</b>	<b>15,3</b>

In the industrial farming, the anti-PRRS antibody seroprevalence was high unlike the low level of postvaccinal anti-erysipelas antibodies. In households, PRRS virus incidence was low and seroconversion after anti-erysipelas vaccination was high (fig. 5).

pulmonary pasteurellosis, colibacillosis, salmonellosis and others (ecchymosis, pulmonary petechia, bronchopneumonia, fibrinous polyserositis, enlarged spleen, large and hemorrhagic lymph nodes, catarrhal and catarrhal-congestive enteritis).



**Figure 5.** Immune response after anti-erysipelas vaccination in households

The pathological and bacteriological examinations carried out on wounds from various organs (lungs, lymph nodes, liver, spleen, intestine) revealed the presence of pathogenic or potentially pathogenic associated bacteria (*Mannheimia (Pasteurella) haemolytica* - 1 strain, *Pasteurella multocida* - 2 strains, *Pseudomonas aeruginosa* - 2 strains, *Streptococcus spp.* - 3 strains, *Klebsiella pneumoniae* - 1 strain, *Escherichia coli* - 14 strains).

The necropsy revealed no characteristic clinical signs, but secondary infection-specific lesions were noticed: Glasser disease,

**Conclusions**

Results from both previous and final research in farms indicate a negative

correlation between the presence of the virus and the postvaccinal immune response.

Therefore, after anti-erysipelas vaccination in pig farms which are distinct by immune status and rearing conditions, seroconversion was low (5.8%, 23.7% and 9%; with a 12.8% mean) following the high PRRS seroprevalence, regardless of the type of anti-erysipelas vaccine used.

In comparison with these results, seroconversion in households was high (57.5%) after anti-erysipelas vaccination, as a result of very low PRRS seroprevalence (2.3%).

After analysis by ELISA of all the samples from farms and households, the high PRRS seroprevalence (54%) induced a low seroconversion (15.3%) post anti-erysipelas vaccination.

The presence of PRRS virus in swine populations proved that it has an immunosuppressive effect demonstrated by the weak immune response capacity after anti-erysipelas vaccination, the existence of associated bacteria and severe disease progress.

PRRS virus is a pathogen that can interfere with the efficacy of vaccination protocols applied in nowadays swine farming systems.

## References

1. **Bărăităreanu, S., Bezman, M., Marinache, S., Cobzariu, D., Bagrinovschi, G., Oțelea, M., Campeanu, M.V., Ivana, S., Daneș, D.** [2010]. Etiology and pathology in porcine respiratory disease complex [PRDC] In finishing pigs. *Lucr. Șt. 53 [12] Medicină Veterinară, Iași, Partea 4.*
2. **Cum M, Yao, J.C., Zhu, L.D., Fang, S.H., Yang, X.J.** [2006]. PRRS seroprevalence in forty five chinese pig herds. *Proceedings of the 19th IPVS Congress, Copenhagen, Denmark, Vol. 2, p 27.*
3. **Herman V., Pascu, C., Costinar, L., Cătană N., Faur B., Văduva, I., Surpat, A., Irimie, S.** [2010]. Investigation of PRRSV Infection in Fattening Farms. *Lucrări Științifice Medicină Veterinară Vol. XLIII[1], Timișoara.*
4. **Nodelijk, G., van Leengoed, L.A., Schoevers, E.J., Kroese, A.H., De Jong, M.C., Wensvoort, G., Verheijden, J.H.** [1997]. Seroprevalence of porcine reproductive and respiratory syndrome virus in Dutch weaning pigs. *Vet Microbiol, 56[1-2]: 21-32.*
5. **Perianu, T.** [2011]. *Tratat de boli infecțioase ale animalelor. Viroze și boli prionice. Sindromul respirator și de reproducție la suine, Vol II, 93-99.*
6. **Reiner, G., Fresen, C., Bronnert, S., Willems, H.** [2009]. Porcine Reproductive and Respiratory Syndrome Virus [PRRSV] infection in wild boars. *Vet Microbiol 136, 3-4, 250.*
7. **Stancu, A., Olariu-Jurca, I., Olariu-Jurca, A., Fluerașu, L.** [2014]. Observations on pathological lesions in young swine PRRS syndrome after weaning. *Lucr Șt Med Vet Vol. XLVII[1], Timișoara.*
8. **Tukelj, M., Valenak, Z.** [2003]. The Role Of Serology In PRRS Control. *4th International Symposium on emerging and re-emerging pig diseases – Rome June 29th – July 2nd, 2003: 125.*
9. **Chiurciu, V., Răduță, M.M.** [2014]. Seroprevalence of some immunosuppressive agents [PRRSV, *M. hyopneumoniae* and PCV2] in wild boars [*Sus scrofa*] in Romania [2013]. *Lucr Șt Med Vet Vol. XLVII [1], 20-25, Timișoara.*
10. **Chiurciu, V., Tudoran, C., Diaconu, L., Stoica, C., Iacob, I., Raduta, M.M.** [2014]. The influence of some microbial immunosuppressive agents on the effectiveness of immunoprophylaxis programs applied in breeding pigs. *Scientific Works C Series LVIII [3], 34-47, București.*
11. **Weigel, R.M., Firkins, L.D., Scherba, G.** [2000]. Prevalence and risk factors for infection with Porcine Reproductive and Respiratory Syndrome Virus [PRRSV] in swine herds in Illinois [USA]. *Vet. Res. 31: 87-88.*